

Cognitive and behaviour change in MND

What you should know


- The word “cognition” refers to the thought processes that include thinking, reasoning, awareness, perception or remembering
- Over 50% of people with MND can experience reduced abilities in thinking and behaviour
- Most people experience relatively mild changes
- A small proportion (5-15%) will show more significant changes and be diagnosed with ‘motor neurone disease with frontotemporal dementia’ or MND/FTD
- Difficulty concentrating, finding the right words for conversations, solving problems and expressing emotions are some of the symptoms of cognitive and behaviour change in MND
- A general practitioner (GP), neurologist or specialist MND clinic can assist in diagnosing changes in cognition and behaviour

Introduction

Past research suggested that motor neurone disease (MND) only affected the nerve cells (neurons) controlling the muscles that enable us to move, speak, breathe and swallow. However, more recent research has found that over 50% of people with MND can experience reduced abilities in thinking and behaviour¹.

When cognitive and behaviour changes occur in MND, it is because there have been changes in specific areas of the brain called the frontal and temporal lobes (shown right)².

Most people experience relatively mild changes. However, approximately 10–15% of patients with ALS show signs of FTD, and may receive a diagnosis of ‘motor neurone disease with frontotemporal dementia’ or MND/FTD³. Often the symptoms of dementia precede the motor symptoms, sometimes by a number of years.



What do changes in cognition and behaviour look like?

When cognitive and behaviour changes are mild, they may not be noticeable or affect daily life. However, when changes are more pronounced, they can have a negative impact on a person’s lifestyle and relationships. Cognitive and behaviour changes can often be a source of confusion and misunderstanding for the person with MND and their family. While the nature of changes in cognition and behaviour will vary from person to person, some of the most common symptoms are listed below.

Changes in cognition can include difficulties with:

- concentration
- thinking quickly
- learning new things
- recalling information from memory
- reasoning and problem solving
- planning and organising
- awareness and insight into MND symptoms

Changes in behaviour or personality can include:

- rigidity – resistance to changing routine or attempting new activities
- impulsivity – doing or saying things without considering consequences
- irritability – experiencing a ‘shorter fuse’ or reacting with disproportionate anger to events
- disinhibition – difficulties in controlling behaviour or making social judgements, such as saying inappropriate things during social events
- apathy – reduced motivation and less interest in previously enjoyed activities
- difficulty understanding and expressing emotion. For example, difficulty in picking up on emotional cues and responding appropriately

Changes in language can include:

- unusual speech patterns, writing or spelling
- difficulty finding the right word in conversation
- taking longer to respond in conversations
- problems in understanding the meaning of words
- using odd words to get a message across
- repeating particular words or phrases over and over ⁴⁻⁷

Who can help?

If you or someone you know might be experiencing these symptoms, it is important to consult with a general practitioner (GP), neurologist or specialist MND clinic. Clinical neuropsychologists can also assist in diagnosing changes in cognition and behaviour, and can help the person and their family to develop practical strategies to manage symptoms. Strategies may include things like reducing distractions, shorter phrases, memos and aids^{8, 9}.

Points to think about

- Many people with MND do not experience cognitive and behaviour change.
- Symptoms might be caused by factors other than changes in the brain (for example, depression or medication effects) so this needs investigating.
- Clinical neuropsychologists and neurologists can assess changes in cognition and behaviour with a view to helping you to develop practical strategies.
- If changes are noticed by the person with MND or others, supports are available to assist with management of these symptoms.

More information

For more information on cognitive and behaviour change in MND please contact your MND Association or relevant health professional.

To find out about motor neurone disease and other fact sheets in this series contact the MND Association in your state or territory
ph. 1800 777 175
or visit www.mndaustralia.org.au

References

1. Strong MJ, Abrahams S, Goldstein LH, Woolley S, McLaughlin P, Snowden J, Mioshi E, Roberts-South A, Benatar M, Hortobágyi T, Rosenfeld J, Silani V, Ince PG, Turner MR 2017, ‘Amyotrophic lateral sclerosis - frontotemporal spectrum disorder (ALS-FTSD): Revised diagnostic criteria’, *Amyotroph Lateral Scler Frontotemporal Degener.*, 18(3-4): 153–174.
2. Strong MJ, Grace GM, Freedman M, Lomen-Hoerth C, Woolley S, Goldstein LH, Murphy J, Shoesmith C, Rosenfeld J, Leigh PN, Buijini L, Ince P, Figlewicz D 2009, ‘Consensus criteria for the diagnosis of frontotemporal cognitive and behavioural syndromes in amyotrophic lateral sclerosis’, *Amyotroph Lateral Sclerosis* 10(3), 131-146.
3. Hobson EV, McDermott CJ 2016, ‘Supportive and symptomatic management of amyotrophic lateral sclerosis’, *Nature Reviews Neurology*, 12(9):526-38
4. Merrilees J, Klapper J, Murphy J, Lomen-Hoerth C, Miller BL 2010, ‘Cognitive and behavioural challenges in caring for patients with frontotemporal dementia and amyotrophic lateral sclerosis’, *Amyotroph Lateral Sclerosis* 11(3), 298-302.
5. Kasper E, Schuster C, Machts J, Bittner D, Vielhaber S, Benecke R, Teipel S and Prudlo J, 2015, ‘Dysexecutive functioning in ALS patients and its clinical implications’, *Amyotroph Lateral Scler Frontotemporal Degener* 16(3-4):160-71.
6. Oliver D 2007, ‘Palliative Care’ in Kiernan MC (Ed). *Motor neurone disease handbook*, MJAbooks, Sydney.
7. Taylor LJ, Brown RG, Tsermentseli S, Al-Chalabi A, Shaw CE, Ellis CM, Leigh PN, Goldstein LH 2012, ‘Is language impairment more common than executive dysfunction in ALS?’ *Journal of Neurology Neurosurgery and Psychiatry* doi:10.1136/jnnp-2012-303526 Epub Oct 2
8. National Institute for Health and Care Excellence (NICE) 2016, ‘Motor neurone disease: assessment and management’, retrieved from <https://www.nice.org.uk/guidance/NG42/chapter/Recommendations#managing-symptoms> 3
9. Caga J, Hsieh S, Lillo P, Dudley K, Mioshi E 2019, ‘The Impact of Cognitive and Behavioral Symptoms on ALS Patients and Their Caregivers’ *Frontiers in Neurology*, 10:192. doi: 10.3389/fneur.2019.00192.